

DOUBLE PARTON SCATTERING PROCESSES IN PP COLLISIONS AND THE SCALE FACTORS.

A. DEL FABBRO

*Dipartimento di Fisica Teorica, Università di Trieste, Strada Costiera 11,
Miramare-Grignano and INFN, Sezione di Trieste, I-34014 Trieste, Italy.*

E-mail: delfabbr@ts.infn.it

The double parton collisions become increasingly important at high energies. The scale factors are the dimensional parameters which characterize these interactions and are the physical observables that can provide a first quantitative information on the correlations in the hadron structure.

1 Double Parton Collisions

Multiple parton interaction processes, where different pairs of partons have an hard scattering in the same hadronic collision, become experimentally feasible at high energies because of the growing flux of partons. The hard partonic interactions occur independently in different points inside the overlap region of the two colliding hadrons and are separated by a distance of the order of the hadron radius. These processes are characterized by two very different scales: the hadronic dimension and the inverse of the hard scale in the partonic subprocesses. Therefore the interference effects between the partonic interactions can be neglected and the expression of the cross sections for these processes have a simple probabilistic interpretation.

In the case of double parton collisions, for the process $pp \rightarrow AB$, the cross section is

$$\sigma_D(AB) = \frac{m}{2} \int dx_1 dx'_1 dx_2 dx'_2 d^2b \times \Gamma(x_1, x_2; b) \hat{\sigma}^A(x_1, x'_1) \hat{\sigma}^B(x_2, x'_2) \Gamma'(x'_1, x'_2; b) \quad (1)$$

The non-perturbative input in this expression is the two-body parton distribution $\Gamma(x_1, x_2; b)$ where x_1 and x_2 are the momentum fractions and b is the distance of the two partons in transverse space. $\hat{\sigma}^A$ and $\hat{\sigma}^B$ are the partonic cross sections for the production of the final states A and B respectively and $m = 1(m = 2)$ when A and B are indistinguishable (distinguishable) processes.

If one assumes that the the momentum fractions are not correlated and the dependence on b can be factorized, the inclusive double parton scattering cross section for the two parton processes A and B reduces to the simplest

expression

$$\sigma_D = m \frac{\sigma_A \sigma_B}{2\sigma_{eff}} \quad (2)$$

where σ_A and σ_B are the inclusive single scattering cross sections for producing the final states A and B respectively. The information on the structure of the hadron in transverse space is summarized in the value of the scale factor σ_{eff} which is universal i.e. does not depend on the considered process. The experimental value quoted by CDF is $\sigma_{eff} = 14.5 \pm 1.7_{-2.3}^{+1.7}$ mb¹.

Using this simple factorized expression one find that the double parton collisions give important effects in several processes at the LHC, for example in the production of $(W, Z)b\bar{b}$, $W+jets$, four $jets$, $b\bar{b}b\bar{b}$, etc². An important case is represented by the production of $Wb\bar{b}$ in a double parton scattering, which is a large source of background in the Higgs boson production mechanism $pp \rightarrow W(H \rightarrow b\bar{b})$ ³.

2 Scale Factors

The scale factor σ_{eff} is related to the transverse size of the region where the hard interactions take place. More precisely, this parameter is given by the overlap in transverse space of the matter distribution of the two interacting hadrons. The measured value of the scale factor is sizably smaller than the naive expectations and it is an indication of important correlations effects in the hadron structure⁴. In fact, while correlations in the momentum fractions will affect the two-body parton distributions only for large values of x , the correlations in the transverse parton coordinates are always present due to the binding force. In a recent paper⁵ the small value of the effective cross section was explained by a simple correlated model of the partonic population in the proton. In this model the distributions in transverse space of gluons and sea quarks are correlated with the configuration taken by the valence quarks, in such a way that extended configurations of the valence give rise to a large number of partons while compact configurations give rise to a smaller number, as one expects for the growth of the energy of the gluonic field with the distance between the valence quarks. On the other hand in the model the momentum fractions of the initial parton pairs are uncorrelated. One of the main features is that there are two different kinds of partons as far as the distributions in transverse space are concerned: the valence quarks and the sea quarks and gluons. As a consequence the scale factor in a reaction involving valence quarks will be different than the scale factor in a reaction involving sea quarks and gluons⁶.

The inclusive double parton scattering cross section is therefore expressed as

$$\sigma_D(A, B) = \frac{m}{2} \sum_{ijkl} \int dx_1 dx'_1 dx_2 dx'_2 d^2b \times \Gamma_{ij}(x_1, x_2; b) \hat{\sigma}_{ik}^A(x_1, x'_1) \hat{\sigma}_{jl}^B(x_2, x'_2) \Gamma'_{kl}(x'_1, x'_2; b) \quad (3)$$

where the indices i and j refer to the different kinds of partons, while dependence of Γ on x_1 , x_2 and b is factorized: $\Gamma_{ij}(x_1, x_2; b) = G_i(x_1)G_j(x_2)F_j^i(b)$, and $G_i(x)$ are the usual one-body parton distributions.

The model introduces the geometrical coefficients⁶

$$\Theta_{kl}^{ij} = \int d^2b F_k^i(b) F_l^{j'}(b), \quad (4)$$

which have the dimension of the inverse of a cross section, so that the double scattering cross section is given by

$$\sigma_D(A, B) = \frac{m}{2} \sum_{ijkl} \Theta_{kl}^{ij} \sigma_{ij}(A) \sigma_{kl}(B) \quad (5)$$

where $\sigma_{ij}(A)$ is the inclusive cross section for the process $ij \rightarrow A$.

By using the equation

$$\sum_{ijkl} \Theta_{kl}^{ij} \sigma_{ij}(A) \sigma_{kl}(B) = \frac{\sigma(A) \sigma(B)}{\sigma_{eff}} \quad (6)$$

we can therefore obtain the scale factors, as a function of the c.m. energy, in various reactions where the rate of double parton scatterings is large enough to measure the corresponding scale factor. We have considered the following production processes: equal sign WW ⁷, Wjj , $Wb\bar{b}$, 4-jets and $b\bar{b}b\bar{b}$. Our results are plotted in Fig.1, and show strong difference between the scale factors in the channels with and without W bosons, as a consequence of the different role played by the valence in the two cases. The scale factor in reactions involving valence quarks is in fact larger than that in reactions involving gluons and sea quarks. This is due to the fact that in the model the average transverse distance of a pair of valence quarks is larger than the average transverse distance of a pair of gluons or sea quarks. The overall qualitative indication of our calculations is that the scale factors are process and energy dependent and that they are the physical observables which can provide a first quantitative information on the correlations in transverse space of the initial state partons.




Figure 1. σ_{eff} as a function of the c.m. energy in various processes in pp collisions.

Acknowledgments

This work was partially supported by the Italian Ministry of University and of Scientific and Technological Research by means of the Fondi per la Ricerca scientifica-Università di Trieste.

References

1. F. Abe et al. (CDF Collaboration), *Phys.Rev.Lett.* **79** 584 (1997); *Phys. Rev.* **D56** 3811 (1997).
2. S. Catani et al. CERN-TH-2000-131, hep-ph/0005025.
3. A. Del Fabbro and D. Treleani, *Phys. Rev.* **D61** 077502 (2000).
4. G. Calucci and D. Treleani, *Phys. Rev.* **D57** 503 (1998).
5. G. Calucci and D. Treleani, *Phys. Rev.* **D60** 054023 (1999).
6. A. Del Fabbro and D. Treleani, *Phys. Rev.* **D63** 057901 (2001).
7. A. Kulesza and W.J. Stirling *Phys. Lett.* **B475** 168 (2000)